

205T40" 6222650

Non-Transduced hMSCs in Immuno-competent mouse muscle

Day 3

DiI + hMSCs





FIG. 1α

+


FIG. 1b

Human MSCs in Mouse Muscle: Day 7

(H&E)



DiI labelled cells +
Sybr-green staining
of all nuclei





+

205TH0" 62226660

FIG. 1c

Human MSCs in Mouse Muscle: Day-7

(H&E, 20X)




3 / 19

205140-62226560


FIG. 1d

Human MSCs in Mouse Muscle: Day 14

Fluorescent Field, 10X



H&E 40X




4 / 19

205T4D" 62226660


FIG. 1e

DiI-Positive α -Gala Transduced hMSCs in
Immuno-competent mouse muscle

d29



d43



5 / 19

205740" 62226650


+

6 / 19

FIG. 1f

Human α GalA-Transduced MSCs in Mouse Muscle: Day-57

10X




+

205T40" 6226660

FIG. 2

D3: IM injection of DiI-labeled PBMCs



7 7 19

205140" 6226650

FIG. 3


D3 IM injection of saline



203TH0" 62226660

FIG. 4

Red and Blue Florescence Channels merged



205THD" 6226660

FIG. 5

α GalA Immunostain of Muscle from Mouse
injected IM with α GalA-hMSCs: 14 days




+

11/ 19

FIG. 6

α -GAL activity in Plasma of Mice
IM-injected with α -GALA-Transduced h MSCs



#: No MSCs, *n=2,

Fabry Mice plasma: 0.5 nmole/h/ml, PNAS 97:7515

+


+

12 / 19


FIG. 7α

Human-specific Alu esquence staining of hMSCs injected into Immuno-competent mouse muscle

Dil-positive cells



Alu-sequence positive cells




+

205THO" 62226660

+

FIG. 7b

Human MSCs in Mouse Muscle: 43 days



137 19

+

205THO" 6226660

Day 57 DiI/Alu






FIG. 7c

14 / 19

+

F1 G. 8 α

α Gala Activity in Muscles of Fabry-KO mice:
14 days after IM injection of α Gala-Transduced hMSCs



Tx = α Gala Transduced cells injected into muscle


Con = Vehicle alone injected into muscle

+

+

16 / 19

FIG. 8b




+

205THD-62226660

FIG. 9a


α Gala Immunostain of Liver from KO Mouse
injected IP with α Gala-hMSCs on Cultispheres:
14 days



IP1Gall-20x

FIG. 9b

α GalA Immunostain of Liver from KO Mouse
injected IP with α GalA-hMSCs on Cultispheres:
14 days




IP2Galla-20x

+

19 / 19

FIG. 10

Intra-Peritoneal Delivery of α Gala-Transduced hMSCs
to Fabry Knock-Out Mice



Increase in α Gala is 5% of normal mouse liver activity (41 nmole/mg)

+